

Review

Chinese Medicine: A Hope for Neurodegenerative Diseases?

Betty Yuen Kwan Law^{a,b}, An Guo Wu^a, Min Jun Wang^{a,b} and Yi Zhun Zhu^{a,b,*}

^a*State Key Laboratory of Quality Research in Chinese Medicine, Macau University of Science and Technology, Macau, China*

^b*School of Pharmacy, Macau University of Science and Technology, Macau, China*

Accepted 12 May 2017

Abstract. With the increase in the proportion of aged population due to the rapid increase of life expectancy, the worldwide prevalence rate of multiple neurodegenerative diseases including Alzheimer's disease, Parkinson's disease, and Huntington's disease has been increased dramatically. The demographic trend toward an older population has drawn the attention to new drug discovery and treatment on age-related diseases. Although a panel of drugs and/or therapies are currently available for treating the neurodegenerative diseases, side effects or insufficient drug efficacy have been reported. With the long history in prescription of Chinese medicine or natural compounds for modulating aged-related diseases, emerging evidence was reported to support the pharmacological role of Chinese medicine in ameliorating the symptoms, or interfering with the pathogenesis of several neurodegenerative diseases. This review brings evidence about today's trends and development of a list of potential neuroprotective herbal compounds from both the traditional and modern pharmacological point of view. With future projections, the potential hope and implication of using Chinese medicine as an alternative source for novel drug discovery for neurodegenerative diseases is proposed.

Keywords: Aging, α -synuclein, Alzheimer's disease, amyloid- β , autophagy, Chinese medicine, disease proteins, leonurine, neurodegenerative diseases, Parkinson's disease

THE AGING WORLD: PATHOGENESIS AND CURRENT THERAPY OF NEURODEGENERATIVE DISEASES

Based on the "World Population Ageing Report 2015" issued by the United Nations, it was postulated that the number of people in the world with age over 60 will accelerate from 901 million to 2.1 billion by 2050 [1]. The population of China is also aging. It was reported that by 2030, 358 million of population will be aged over 60, and younger generations will have a heavy burden of elderly care. Consistently,

worldwide data have indicated an increasing prevalence of neurodegenerative diseases including Alzheimer's disease (AD), Parkinson's disease (PD), and less common neurodegenerative disease such as Huntington's disease (HD) with age [2], which is worth our attention. It is still a big challenge for academia and pharmaceutical industries to discover a really powerful anti-neurodegenerative agent based on the high failure rate of the AD candidate compounds in the last two decades.

Alzheimer's disease

Dementia is considered as one of the most important health problems in elderly, and AD has accounted for about 60% of all dementias among elderly in

*Correspondence to: Prof. Yi-Zhun Zhu, Room N401, Block N, School of Pharmacy, Macau University of Science and Technology, Avenida Wai Long, Taipa, Macau, China. Tel.: +853 8897 2181; E-mail: yzzhu@must.edu.mo.

the world [3]. In the US, there are approximately 5 million of AD patients currently, and it is postulated to have more than 13 million of AD patients by 2050. The costs of care to AD patients is predicted to reach a huge amount of \$1.2 trillion in the US by 2050 [4]. The systematic epidemiology analysis by Chan et al. reported the number of AD patients in China from 1999 to 2010 has been increased from 1.93 to 5.69 million [5]. AD is characterized by progressive loss of memory, cognitive and decision making deficits with problems in language ability. These symptoms were accompanied by various neuropsychiatric conditions such as anxiety, depression, or apathy [6]. Senile plaques containing amyloid- β ($A\beta$) peptide produced by sequential cleavage of the amyloid- β protein precursor ($A\beta$ PP) via β - and γ -secretases [7], and neurofibrillary tangles (hyperphosphorylated microtubule-associated tau protein) in the neocortex, are highly correlated with neuropathological features and neuronal loss of AD. Formation of plaque and tangles is potentiated by various factors such as genetic, inflammation, and production of reactive oxygen species (ROS) [8]. The amyloid cascade hypothesis suggested the production of toxic $A\beta$ aggregates lead to the pathogenesis of AD. The discovery of mutations in presenilin (PS1), which is involved in $A\beta$ PP cleavage and $A\beta$ production [9], has further supported the hypothesis. Although the amyloid hypothesis is widely accepted, with the recent failure of a clinical study of $A\beta$ vaccination, new therapeutic strategy for AD, such as the use of immunotherapies with the antibodies or tau-targeted therapies, are currently under investigation [10].

Parkinson's disease

With a prevalence of over 1,900 patients per 100,000 in people of age over 80, PD is characterized by the death of dopaminergic neurons in the substantia nigra region and Lewy body deposition in the midbrain. This is accompanied by movement disorder symptoms such as bradykinesia, rigidity, and resting tremor. Pathogenic risk factors of PD include microtrauma or focal cerebrovascular damage of brain, stimulation by toxins, drugs or pesticides, dysfunction of ubiquitin-proteasome system, oxidative stress and neuroinflammatory reactions, genomic factors which lead to conformational changes, misfolding and aggregation of proteins, and neuronal death [11]. Current pharmacological interference for PD includes the use of dopamine precursors such

as levodopa (L-DOPA); dopamine agonists such as amantadine, apomorphine, bromocriptine, cabergoline, pergolide, and ropinirole; monoamine oxidase (MAO) inhibitors such as selegiline; and catechol-O-methyltransferase (COMT) inhibitors such as entacapone [2]. However, motor fluctuations and dyskinesia are the major complications of L-DOPA therapy. Additional complications include gastrointestinal or cardiovascular problems, changes in neuroendocrine system, and psychiatric disorders after treatment of anti-parkinsonian drugs [12].

Huntington's disease

HD is an inherited neurodegenerative disease characterized by motor disorder and cognitive problems in dementia, and is fatal in majority of cases. With an average onset age at around 40, psychiatric problems such as depression or anxiety were also observed in patients of HD. HD is more prevalent in European peoples with appropriately 10 to 15 patients out of 100,000 people. Pathogenesis of HD is caused by a mutation with an over 40 number of CAG trinucleotide repeat in the huntingtin (HTT) gene, resulting an excessively long polyglutamine tract near the N-terminus of HTT protein, which is responsible for early development of neurons [13]. Although currently there is still no effective therapeutic treatment to slow down the decline in motor or cognitive function in HD, symptoms such as chorea can be modulated by dopamine receptor blockers like haloperidol or dopamine-depleting agent such as tetrabenazine [14].

TRADITIONAL CHINESE MEDICINE (TCM) HERBAL PRESCRIPTION AND THEORIES FOR MODULATING AGE-RELATED NEURODEGENERATIVE DISEASES

TCMs has been prescribed in the Chinese community for more than 2,000 y for treating symptoms including forgetfulness, disorientation, insomnia, loss of consciousness, cramps, and seizures, which are highly related to aging or age-related diseases [15]. According to Chinese medicine theory, age-related neurodegeneration are attributed to the follow aspects: 1) The loss of kidney function (shen jing) is accompanied by a lack of nutrients to the brain, which leads to the gradual diminishing of the brain and causes dementia; 2) Heart and spleen function deficiency and a lack of blood (qi) flow that reaches

the brain, leads to symptoms such as nervousness, palpitations, insomnia, anxiety, and a decrease in learning and memory ability; 3) The stagnation and stasis of blood (qi) in the peripheral organs lead to insufficient blood flow to the brain and causes the decline in memory [16, 17]. Currently, the top 10 popular anti-aging herbs are Heshouwu (Fleece-Flower Root) [18], Huangqi (*Radix Astragalii*) [19], Renshen (*Radix Ginseng*) [20], Sanqi (*Radix Notoginseng*) [21], Ciwujia (*Radix Acanthopanaxis Senticosli*) [22], Lingzhi (*Ganoderma*) [23], Gouqi (*Fructus Lycii*) [24], Hongjingtian (*Radix et Rhizoma Rhodiola kirilowii*) [25], Jiaogulan (*Herba Gynostematis Pentaphylli*) [26], and Fengwangjiang (Royal Jelly) [27] in China [28]. Most of them are traditional TCMs prescribed for maintaining good memory and balanced body homeostasis, regulating age-related neurodegenerative diseases, or promoting longevity among Chinese people. Modern pharmacological researchers have confirmed part of their beneficial roles in term of anti-oxidation [21], anti-inflammation [29, 30], and anti-apoptosis properties [19].

As neurodegeneration is a complicated and multifactorial disease, Chinese medicine offers the advantage of modulating the systemic organs via synergetic effects of different compositions within a single herb. For example, the well-known TCMs such as Yuanzhi (*Radix Polygalae*) [31], Renshen (*Radix Ginseng*) [32], and Hehuanpi (*Cortex Albiziae*) [33] are prescribed to ameliorate memory loss; Gouqi (*Fructus Lycii*), Fuling (*Poria*), Gancao (*Radix Glycyrrhizae*), and Shichangpu (*Rhizoma Acori Tatarinowii*) are used to improve the cognitive function [34]. In addition, traditional herbal formulations such as “Kaixinsan” which are composed of 4 herbs, *Radix Ginseng*, *Radix Polygalae*, *Rhizoma Acori Tatarinowii*, and *Poria*, is a famous TCM formula for the treatment of mental disorders, forgetfulness, and depression possibly via the pCREB pathways [35]. Additional herbal formulations such as Danggui-Shaoyao-San, Tianma Gouteng Yin, and Jitai Tablet, are reported to exert their neuroprotective effects via ameliorating cognitive dysfunction in AD model [36], preventing degeneration of dopaminergic neurons in PD model [37], and improving behavioral impairments in PD mice [38], respectively. *Ginkgo biloba* extract EGb 761[®], and a modified formulation of SuHeXiang Wan, were found effective in modulating cognitive behavior and neuropsychiatric symptoms of AD [39], A β -induced memory impairment, and A β levels in AD mice model [40],

respectively. Furthermore, our previous work on *Herbal Leonuri* extract [41] has confirmed the cerebral protection effect of the herbal extract on stroke model via its anti-oxidation properties. Therefore, the Chinese medicine provides a rich source for the identification and pharmacological research of potential neuroprotective compounds for therapeutic application.

MODERN PHARMACOLOGICAL ROLE OF TCMS AND THEIR MOLECULAR MECHANISMS IN REGULATING NEURODEGENERATIVE DISEASES

Autophagic degradation of neurodegenerative disease proteins

Autophagy is a cellular system responsible for delivering cellular protein, organelles, or invading microorganisms to lysosomal degradation. With a variety of physiological and pathophysiological roles such as starvation adaptation or clearance of cytoplasmic constituents extensively reported in the past decades, the pathological role of autophagy in cancer, aging, and neuro-degenerative disorders have also been well studied [42]. It was reported that the ubiquitin-proteasomal system is responsible for cellular degradation of unwanted or excessive proteins and organelles [43]; however, membrane-bound proteins, aggregated, and oligomeric neurodegenerative disease proteins, which are too large to pass through the narrow barrel of the proteasome, are degraded by autophagy [44]. Examples of these proteins include HTT protein with long mutant polyglutamine (polyQ) tracts [45, 46], and A53T/A30P mutant α -synuclein protein, which contributes to the pathogenesis of HD and PD respectively, can be degraded by autophagy [45–51]. Autophagy was reported to play a protective role in alleviating the levels and hence, the toxicity of both mutant huntingtin and α -synuclein [52, 53], suggesting the pharmacological activation of autophagy may be beneficial to regulate neurodegenerative diseases [54].

Concomitantly, recent studies have reported the identification of novel autophagic enhancers from TCMs as potential neuroprotective agents. Our previous work have identified onjisaponin B [55] and a partially purified bioactive fraction from *Radix Polygalae* [56] as novel autophagic activators that are able to accelerate the degradation, and decrease the toxicity of mutant α -synuclein and HTT protein in cellular models. Furthermore, we have reported the

active component, neferine, isolated from the lotus seed embryo of *Nelumbo nucifera*, can also attenuate the protein level and toxicity of mutant HTT through autophagy [57]. Hederagenin and α -hederin isolated from *Hedera helix* are reported to activate autophagy, promote degradation of neurodegenerative diseases proteins, and improve motor deficits in mice treated with neurotoxin, MPTP [58]. Recently identified TCMs, such as arctigenin from *Arctium lappa* (L.), are able to accelerate the autophagic clearance of A β [59]; isorhynchophylline, an alkaloid from *Uncaria rhynchophylla*, facilitates the autophagic degradation of α -synuclein in neuronal cells [60], with improvement in cognitive deficit of A β - rats [61]; neuroprotective effects of baicalein from *Scutellaria baicalensis* [62], and resveratrol from red wine or food [63] were also suggested to be executed possibly via autophagy. All of the above findings have suggested the potential therapeutic roles and new directions for isolating novel autophagic activators from TCMs for neuro-therapy.

Novel herbal compounds with neuroprotective effects via regulating different signaling pathways

Our previous findings suggested that leonurine, an alkaloid isolated from *Herba leonuri* (HL), exerts its neuroprotective effect against cerebral ischemia/reperfusion-induced mitochondrial dysfunctions in middle cerebral artery occlusion model of rat brain, via its anti-oxidation and anti-apoptosis properties [41]. Besides, leonurine decreased the infarct volume

and further improved neurological deficit of stroke rats. The protective effects on ischemic stroke models were confirmed to be executed via an increase in the activities of anti-oxidation enzymes, such as glutathione peroxidase and superoxide dismutase, with the decrease in mitochondrial ROS production and adenosine triphosphate (ATP) biosynthesis [64]. In addition, SCM198 (4-guanidino-n-butyl syringate), a new compound from HL, was proved to attenuate the neurotoxin (6-OHDA)-induced apoptotic cell death in dopaminergic cells and improve rotational behavior in 6-OHDA-induced motor deficit rat model via an increase in superoxide dismutase antioxidant activity with a decrease in levels of ROS [65]. Furthermore, SCM-198 was confirmed to ameliorate the recognition ability, spatial memory, cognitive deficits, and promote survival of neurons in A β PP and PS1 transgenic mice [66], suggesting the potential therapeutic value of TCMs on neuroprotection (Fig. 1).

Salidroside isolated from *Rhodiola sachalinensis* protects cells against cytotoxicity and oxidative damage induced by accumulation of A β [67, 68]. Furthermore, salidroside possess its neuroprotective effect via improving the impaired hippocampal neurogenesis by scavenging ROS in the streptozotocin induced AD rat model [69]. Curcumin, a well-known natural product with potent anti-inflammation and anti-oxidation effects, was reported to bind plaques, reduce and inhibit the formation A β oligomers and fibrils in mice model [70]. Huperzine A, an alkaloid extracted from the plant *Huperzia serrate*, is commonly used in China to treat AD [71]. Clinical trials have demonstrated the protective effect of




Fig. 1. SCM-198, derived from *Herba leonuri*, shows neuroprotective potential in both 6-OHDA induced PD mice and A β PP/PS1 double-transgenic AD mice.

huperzine A in improving memory deficits in patients with senescent forgetfulness symptoms [72]. It was reported that huperzine A can rescue neural stem cells from apoptosis induced by A β peptide [73]. Besides, huperzine A can work as a potential therapeutic agent for AD via inhibiting acetylcholinesterase in primary cultures and reduces neuronal toxicity induced by glutamate [74].

Recently, the neuroprotective effects of natural product, garlic (*Allium sativum*) extract were studied. For example, *Allium sativum* extract co-cultivated with *Hericium erinaceus mycelium* showed neuroprotective effects in ischemic *in vivo* model [75]. Treatment of garlic extract and methionine can attenuate toxic effects induced by neurotoxicant mercury possibly via their anti-oxidation properties [76]. *Allium* vegetable is one of the rich sources for organic sulfur-containing compounds. It has been proven that diallyl disulfide, diallyl trisulfide, and S-allyl cysteine, which are the key ingredients of garlic, are hydrogen sulfide donors. In fact, the lack of hydrogen sulfide contributes to pathogenesis of arterial and pulmonary hypertension, AD, liver cirrhosis, and gastric mucosal injury [77]. Therefore, identification of organic molecules that can regulate the formation of hydrogen sulfide is important. Concomitantly, our group reported that S-propargyl cysteine (ZYZ-802), a novel hydrogen sulfide-modulated agent, can modulate ischemic heart disease [77], and improve spatial learning and memory deficits induced by lipopolysaccharide-mediated neuro-inflammation through regulating the hydrogen sulfide pathway [78]. All these have suggested the neuroprotective role of hydrogen sulfide-modulated agent present in natural product or TCMs.

PERSPECTIVE: THE USE OF TRADITIONAL TCMS FOR MODULATING NEURODEGENERATIVE DISEASES AND CHALLENGES

Current therapeutic approaches and challenges for neurodegenerative diseases

With the continuous extension of human life expectancy followed by an increasing number of patients with dementia (especially AD), researchers focus not only on the pathogenesis of AD, but also on the discovery of drugs for prevention or treatment of the disease. However, up to now, there is still no effective treatment for AD, and only treatments attempting to counterbalance the disturbance by

neurotransmitters are available for relieving the symptoms currently. For example, 3 cholinesterase inhibitors, donepezil, rivastigmine, and galantamine are approved drugs for the treatment of mild to moderate AD now [79, 80]. Besides, anti-psychotic and anti-depressant drugs such as serotonin reuptake inhibitors (fluoxetine and fluvoxamine) [81] are available for alleviating the behavioral symptoms of dementia [82].

Other than the symptomatic approach, treatments that aim to regulate the pathogenesis of the disease are under extensive investigation. These ‘disease-modifying’ drugs, which focus on interfering the progression of disease at different levels, for example, modulation of deposition of A β plaques and formation of neurofibrillary tangle or tau protein, are under clinical trials or experimental validation. For example, A β lowering agents such as β -secretase enzyme BACE1 inhibitor (CTS-21166) and γ -secretase inhibitor (LY-450139), are under Phase I and Phase III clinical trials, respectively. However, inhibition of BACE1 led to toxic side effect due to the other essential physiological roles of BACE1. Besides, due to the large size of the active site of BACE1, compounds with binding affinity to BACE are unlikely to cross the blood brain barrier [6]. Similarly, other than A β , γ -secretase can also cleave other transmembrane proteins which are essential for growth and development, such as notch receptor 1, and therefore affect the development of BACE1 inhibitors or γ -secretase inhibitors as potential therapeutic agents of AD [83]. Drugs interfering with tau deposition (methylene blue, phenothiazine, or methylthioninium chloride) [84] or tau phosphorylation (pyrazolopyrazines, lithium, pyrazolopyridines, aminothiazole AR-A014418, or sodium valproate) [85] are under clinical trial evaluation. Other approaches, such as the use of anti-inflammatory (NSAIDs), or oxidation drugs, such as vitamin E, Ginkgo biloba, natural polyphenols including blueberries, green tea, curcumin, ω 3 fatty acids, red wine, folate, vitamin B6, and vitamin B12, are also under intensive investigation [6].

Up to now, levodopa remains the most effective therapy for alleviating the symptoms of PD; however, severe side effects such as “wearing off” effect, dyskinesia, fluctuations, and other motor deficits remained unresolved [86]. Other than pharmacological approaches, neurosurgical treatment via deep brain stimulation is an alternative treatment of PD [87]. Although many management strategies have been made with progress in enhancing the effect of

PD therapy with minimum side effects, in addition to modulate the disease by the symptomatic approach, advance treatments targeting the pathogenesis of the disease are highly desirable.

The use of TCMs as new therapeutic approach for modulating aged-related neurodegenerative diseases

This report has highlighted the discovery of new chemicals from TCMs or natural products for modulating age-related neurodegenerative diseases from different approaches (Fig. 2), for example, accelerating the clearance of neurodegenerative disease proteins via induction of autophagy. Rapamycin, a well-known mTOR inhibitor, is effective in inducing the clearance of mutant proteins via autophagy [53]. However, mTOR inhibitors possess side effects by inhibiting immunological function, cellular protein synthesis, and proliferation; therefore, identification of novel compounds that activate the mTOR-independent induction of autophagy, has become an alternative approach in neuro-therapeutic drug discovery. For example, lithium which is naturally

present in grains and vegetables, can activate autophagy via inhibition of inositol monophosphatase, leading to depletion of free inositol and decrease in myo-inositol-1,4,5-triphosphate (IP₃) levels [48], together with a significant decrease in mutant huntingtin and α -synuclein level, suggesting the therapeutic potential of autophagic enhancer in neuro-therapy.

Although the use of TCMs may serve as an alternative way to prevent or treat aged-related neurodegenerative diseases, incorrect dosage, unknown molecular targets and mechanisms, or high toxicity of herbal decoction due to its complexity of compounds formulation remain the major obstacles in developing TCMs for therapeutic uses [88]. However, with a long history in prescribing TCMs for preventing or modulating or age-related disorders, dementia, or its symptoms in the Chinese local community [89, 90], interlinking modern neuropharmacology with therapeutic effect of TCMs has been a highlighted research field in these past decades. By combining advanced isolation and accurate characterization analysis on the chemical compositions of herbal formula, for example, by using ultra-performance




Fig. 2. Schematic diagram showing the modulation of age-related neurodegenerative diseases via the anti-inflammation, anti-oxidation, anti-apoptosis, and autophagy effects exerted by the listed popular TCMs in the Chinese local community. By combining the traditional therapeutic effects of the herbal compounds reported in ancient literature with modern pharmacological mechanisms reported in scientific literature, the hope of developing TCMs as potential neuroprotective agents is highlighted.

liquid chromatography coupled to (quadrupole) time-of-flight mass spectrometry UHPLC-(Q)TOF-MS [56], together with modern biotechnology including computation docking [91], proteomics [92], and other advanced technology such as molecular bio-probe labeling on herbal compounds, live cell imaging, or genetic knockout and RNA interference technology [93], all lead to breakthroughs in the pharmacological research in TCM study and provide us with hope for developing TCMs as alternative medicine for modulating neurodegenerative diseases.

ACKNOWLEDGMENTS

This work was supported by the FDCT grant from the Science and Technology Development Fund of Macao (Project code: 090/2013/A3, 039/2016/A and 055/2016/A2), and Faculty Research Grant of MUST (FRG-17-006-SP).

Authors' disclosures available online (<http://j-alz.com/manuscript-disclosures/17-0374r1>).

REFERENCES

- [1] World Population Ageing 2015, http://www.un.org/en/development/desa/population/publications/ageing/WPA2015_Infochart.shtml
- [2] Cacabelos R (2017) Parkinson's disease: From pathogenesis to pharmacogenomics. *Int J Mol Sci* **18**, 551.
- [3] Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, Hall K, Hasegawa K, Hendrie H, Huang Y, Jorm A, Mathers C, Menezes PR, Rimmer E, Sczufca M, Alzheimer's Disease International (2005) Global prevalence of dementia: A Delphi consensus study. *Lancet* **366**, 2112-2117.
- [4] Sibener L, Zaganjor I, Snyder HM, Bain LJ, Egge R, Carrillo MC (2014) Alzheimer's Disease prevalence, costs, and prevention for military personnel and veterans. *Alzheimers Dement* **10**, S105-S110.
- [5] Chan KY, Wang W, Wu JJ, Liu L, Theodoratou E, Car J, Middleton L, Russ TC, Deary IJ, Campbell H, Wang W, Rudan I, Global Health Epidemiology Reference Group (GHERG) (2013) Epidemiology of Alzheimer's disease and other forms of dementia in China, 1990-2010: A systematic review and analysis. *Lancet* **381**, 2016-2023.
- [6] Yiannopoulou KG, Papageorgiou SG (2013) Current and future treatments for Alzheimer's disease. *Ther Adv Neurol Disord* **6**, 19-33.
- [7] Zhang YW, Xu H (2007) Molecular and cellular mechanisms for Alzheimer's disease: Understanding APP metabolism. *Curr Mol Med* **7**, 687-696.
- [8] Govaerts L, Schoenen J, Bouhy D (2007) [Pathogenesis of Alzheimer's disease: Molecular and cellular mechanisms]. *Rev Med Liege* **62**, 209-216.
- [9] De Strooper B, Saftig P, Craessaerts K, Vanderstichele H, Guhde G, Annaert W, Von Figura K, Van Leuven F (1998) Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* **391**, 387-390.
- [10] Hasegawa M (2016) Molecular mechanisms in the pathogenesis of Alzheimer's disease and tauopathies-prion-like seeded aggregation and phosphorylation. *Biomolecules* **6**, 24.
- [11] Tramutola A, Lanzillotta C, Perluigi M, Butterfield DA (2016) Oxidative stress, protein modification and Alzheimer disease. *Brain Res Bull*. doi: 10.1016/j.brainresbull.2016.06.005
- [12] Owolabi LF, Samaila AA, Sunmonu T (2014) Gastrointestinal complications in newly diagnosed Parkinson's disease: A case-control study. *Trop Gastroenterol* **35**, 227-231.
- [13] Dayalu P, Albin RL (2015) Huntington disease: Pathogenesis and treatment. *Neurol Clin* **33**, 101-114.
- [14] Huntington Study Group (2006) Tetrabenazine as anti-chorea therapy in Huntington disease: A randomized controlled trial. *Neurology* **66**, 366-372.
- [15] Wu TY, Chen CP, Jinn TR (2011) Traditional Chinese medicines and Alzheimer's disease. *Taiwan J Obstet Gynecol* **50**, 131-135.
- [16] Hempen CH, Fischer T (2009) *A Materia Medica for Chinese Medicine: Plants, Minerals and Animal Products*, Elsevier Health Sciences, London, UK.
- [17] Law BY, Mok SW, Wu AG, Lam CW, Yu MX, Wong VK (2016) New potential pharmacological functions of Chinese herbal medicines via regulation of autophagy. *Molecules* **21**, 359.
- [18] Yu J, Xie J, Mao XJ, Wei H, Zhao SL, Ma YG, Li N, Zhao RH (2012) Comparison of laxative and antioxidant activities of raw, processed and fermented Polygoni Multiflori radix. *Chin J Nat Med* **10**, 63-67.
- [19] Sun GG, Shih JH, Chiou SH, Hong CJ, Lu SW, Pao LH (2016) Chinese herbal medicines promote hippocampal neuroproliferation, reduce stress hormone levels, inhibit apoptosis, and improve behavior in chronically stressed mice. *J Ethnopharmacol* **193**, 159-168.
- [20] Li H, Kang T, Qi B, Kong L, Jiao Y, Cao Y, Zhang J, Yang J (2016) Neuroprotective effects of ginseng protein on PI3K/Akt signaling pathway in the hippocampus of D-galactose/AlCl₃ inducing rats model of Alzheimer's disease. *J Ethnopharmacol* **179**, 162-169.
- [21] Huang JL, Jing X, Tian X, Qin MC, Xu ZH, Wu DP, Zhong ZG (2017) Neuroprotective properties of Panax notoginseng saponins via preventing oxidative stress injury in SAMP8 mice. *Evid Based Complement Alternat Med* **2017**, 8713561.
- [22] Li XZ, Zhang SN, Wang KX, Liu SM, Lu F (2014) iTRAQ-based quantitative proteomics study on the neuroprotective effects of extract of *Acanthopanax senticosus* harm on SH-SY5Y cells overexpressing A53T mutant α -synuclein. *Neurochem Int* **72**, 37-47.
- [23] Qiu J, Wang X, Song C (2016) Neuroprotective and antioxidant lanostanoid triterpenes from the fruiting bodies of *Ganoderma atrum*. *Fitoterapia* **109**, 75-79.
- [24] Fang F, Peng T, Yang S, Wang W, Zhang Y, Li H (2016) Lycium barbarum polysaccharide attenuates the cytotoxicity of mutant huntingtin and increases the activity of AKT. *Int J Dev Neurosci* **52**, 66-74.
- [25] Zhang B, Li Q, Chu X, Sun S, Chen S (2016) Salidroside reduces tau hyperphosphorylation via up-regulating GSK-3 β phosphorylation in a tau transgenic *Drosophila* model of Alzheimer's disease. *Transl Neurodegener* **5**, 21.
- [26] Jia D, Rao C, Xue S, Lei J (2015) Purification, characterization and neuroprotective effects of a polysaccharide

- from *Gynostemma pentaphyllum*. *Carbohydr Polym* **122**, 93-100.
- [27] Zamani Z, Reisi P, Alaei H, Pilehvarian AA (2012) Effect of Royal Jelly on spatial learning and memory in rat model of streptozotocin-induced sporadic Alzheimer's disease. *Adv Biomed Res* **1**, 26.
- [28] Top 10 anti-aging herbs, <http://kaleidoscope.cultural-china.com/en/7Kaleidoscope10712.html>, Accessed March 21.
- [29] Nardi GM, Farias Januario AG, Freire CG, Megiolaro F, Schneider K, Perazzoli MR, Do Nascimento SR, Gon AC, Mariano LN, Wagner G, Niero R, Locatelli C (2016) Anti-inflammatory activity of berry fruits in mice model of inflammation is based on oxidative stress modulation. *Pharmacognosy Res* **8**, S42-S49.
- [30] Mok S, Wong V, Law B (2016) Comment: New potential pharmacological functions of Chinese herbal medicines via regulation of autophagy: The search of reliable pharmaceutical candidates for chronic disorders therapy. *J Neurol Neuromed* **1**, 6-9.
- [31] Zhang H, Han T, Zhang L, Yu CH, Wan DG, Rahman K, Qin LP, Peng C (2008) Effects of tenuifolin extracted from radix polygalae on learning and memory: A behavioral and biochemical study on aged and amnesic mice. *Phytomedicine* **15**, 587-594.
- [32] Wan L, Cheng Y, Luo Z, Guo H, Zhao W, Gu Q, Yang X, Xu J, Bei W, Guo J (2015) Neuroprotection, learning and memory improvement of a standardized extract from Renshen Shouwu against neuronal injury and vascular dementia in rats with brain ischemia. *J Ethnopharmacol* **165**, 118-126.
- [33] Dhanya K, Satish S, Karunakar H, Shabaraya AR (2016) Investigation on learning and memory enhancing activity of essential oil in *Albizia julibrissin* flowers in experimental mice. *Asian J Biomed Pharmaceut Sci* **6**, 11-15.
- [34] Liu P, Kong M, Liu S, Chen G, Wang P (2013) Effect of reinforcing kidney-essence, removing phlegm, and promoting mental therapy on treating Alzheimer's disease. *J Tradit Chin Med* **33**, 449-454.
- [35] Dong XZ, Wang DX, Yu BY, Liu P, Hu Y (2016) Kai-Xin-San, a traditional Chinese medicine formulation, exerts antidepressive and neuroprotective effects by promoting pCREB upstream pathways. *Exp Ther Med* **12**, 3308-3314.
- [36] Fu X, Wang Q, Wang Z, Kuang H, Jiang P (2015) Danggui-Shaoyao-San: New hope for Alzheimer's disease. *Aging Dis* **7**, 502-513.
- [37] Liu LF, Song JX, Lu JH, Huang YY, Zeng Y, Chen LL, Durairajan SS, Han QB, Li M (2015) Tianma Gouteng Yin, a Traditional Chinese Medicine decoction, exerts neuroprotective effects in animal and cellular models of Parkinson's disease. *Sci Rep* **5**, 16862.
- [38] Liu J, Gao J, Tu S, Xu S, Liu Y, Shang W, Gu C, Huang Y, Han M (2014) Neuroprotective effects of Jitai Tablet, a Traditional Chinese Medicine, on the MPTP-induced acute model of Parkinson's Disease: Involvement of the dopamine system. *Evid Based Complement Alternat Med* **2014**, 542383.
- [39] Ihl R, Tribanek M, Bachinskaya N, GOTADAY Study Group (2012) Efficacy and tolerability of a once daily formulation of Ginkgo biloba extract EGb 761® in Alzheimer's disease and vascular dementia: Results from a randomised controlled trial. *Pharmacopsychiatry* **45**, 41-46.
- [40] Jeon S, Bose S, Hur J, Jun K, Kim YK, Cho KS, Koo BS (2011) A modified formulation of Chinese traditional medicine improves memory impairment and reduces A β level in the Tg-APPswe/PS1dE9 mouse model of Alzheimer's disease. *J Ethnopharmacol* **137**, 783-789.
- [41] Qi J, Hong ZY, Xin H, Zhu YZ (2010) Neuroprotective effects of leonurine on ischemia/reperfusion-induced mitochondrial dysfunctions in rat cerebral cortex. *Biol Pharm Bull* **33**, 1958-1964.
- [42] Mizushima N (2007) Autophagy: Process and function. *Genes Dev* **21**, 2861-2873.
- [43] Law BY, Wang M, Ma DL, Al-Mousa F, Michelangeli F, Cheng SH, Ng MH, To KF, Mok AY, Ko RY, Lam SK, Chen F, Che CM, Chiu P, Ko BC (2010) Alisol B, a novel inhibitor of the sarcoplasmic/endoplasmic reticulum Ca(2+) ATPase pump, induces autophagy, endoplasmic reticulum stress, and apoptosis. *Mol Cancer Ther* **9**, 718-730.
- [44] Rubinsztein DC (2006) The roles of intracellular protein-degradation pathways in neurodegeneration. *Nature* **443**, 780-786.
- [45] Ravikumar B, Duden R, Rubinsztein DC (2002) Aggregate-prone proteins with polyglutamine and polyalanine expansions are degraded by autophagy. *Hum Mol Genet* **11**, 1107-1117.
- [46] Ravikumar B, Vacher C, Berger Z, Davies JE, Luo S, Oroz LG, Scaravilli F, Easton DF, Duden R, O'Kane CJ, Rubinsztein DC (2004) Inhibition of mTOR induces autophagy and reduces toxicity of polyglutamine expansions in fly and mouse models of Huntington disease. *Nat Genet* **36**, 585-595.
- [47] Webb JL, Ravikumar B, Atkins J, Skepper JN, Rubinsztein DC (2003) Alpha-Synuclein is degraded by both autophagy and the proteasome. *J Biol Chem* **278**, 25009-25013.
- [48] Sarkar S, Floto RA, Berger Z, Imarisio S, Cordenier A, Pasco M, Cook LJ, Rubinsztein DC (2005) Lithium induces autophagy by inhibiting inositol monophosphatase. *J Cell Biol* **170**, 1101-1111.
- [49] Sarkar S, Davies JE, Huang Z, Tunnacliffe A, Rubinsztein DC (2007) Trehalose, a novel mTOR-independent autophagy enhancer, accelerates the clearance of mutant huntingtin and alpha-synuclein. *J Biol Chem* **282**, 5641-5652.
- [50] Berger Z, Ravikumar B, Menzies FM, Oroz LG, Underwood BR, Pangalos MN, Schmitt I, Wullner U, Evert BO, O'Kane CJ, Rubinsztein DC (2006) Rapamycin alleviates toxicity of different aggregate-prone proteins. *Hum Mol Genet* **15**, 433-442.
- [51] Wong E, Cuervo AM (2010) Autophagy gone awry in neurodegenerative diseases. *Nat Neurosci* **13**, 805-811.
- [52] Rubinsztein DC, DiFiglia M, Heintz N, Nixon RA, Qin ZH, Ravikumar B, Stefanis L, Tolkovsky A (2005) Autophagy and its possible roles in nervous system diseases, damage and repair. *Autophagy* **1**, 11-22.
- [53] Rubinsztein DC, Gestwicki JE, Murphy LO, Klionsky DJ (2007) Potential therapeutic applications of autophagy. *Nat Rev Drug Discov* **6**, 304-312.
- [54] Wong VK, Zeng W, Law BY (2017) Perspective: Role of autophagy in neuroprotective properties of traditional Chinese medicines. *J Neuroinfect Dis* **8**, 239.
- [55] Wu AG, Wong VK, Xu SW, Chan WK, Ng CI, Liu L, Law BY (2013) Onjisaponin B derived from Radix Polygalae enhances autophagy and accelerates the degradation of mutant α -synuclein and huntingtin in PC-12 cells. *Int J Mol Sci* **14**, 22618-22641.
- [56] Wu AG, Wong VK, Zeng W, Liu L, Law BY (2015) Identification of novel autophagic Radix Polygalae fraction by

- cell membrane chromatography and UHPLC-(Q)TOF-MS for degradation of neurodegenerative disease proteins. *Sci Rep* **5**, 17199.
- [57] Wong VK, Wu AG, Wang JR, Liu L, Law BY (2015) Neferine attenuates the protein level and toxicity of mutant huntingtin in PC-12 cells via induction of autophagy. *Molecules* **20**, 3496-3514.
- [58] Wu AG, Zeng W, Wong VK, Zhu YZ, Lo AC, Liu L, Law BY (2017) Hederagenin and α -hederin promote degradation of proteins in neurodegenerative diseases and improve motor deficits in MPTP-mice. *Pharmacol Res* **115**, 25-44.
- [59] Zhu Z, Yan J, Jiang W, Yao XG, Chen J, Chen L, Li C, Hu L, Jiang H, Shen X (2013) Arctigenin effectively ameliorates memory impairment in Alzheimer's disease model mice targeting both β -amyloid production and clearance. *J Neurosci* **33**, 13138-13149.
- [60] Lu JH, Tan JQ, Durairajan SS, Liu LF, Zhang ZH, Ma L, Shen HM, Chan HY, Li M (2012) Isorhynchophylline, a natural alkaloid, promotes the degradation of alpha-synuclein in neuronal cells via inducing autophagy. *Autophagy* **8**, 98-108.
- [61] Xian YF, Su ZR, Chen JN, Lai XP, Mao QQ, Cheng CH, Ip SP, Lin ZX (2014) Isorhynchophylline improves learning and memory impairments induced by D-galactose in mice. *Neurochem Int* **76**, 42-49.
- [62] Lu JH, Ardah MT, Durairajan SS, Liu LF, Xie LX, Fong WF, Hasan MY, Huang JD, El-Agnaf OM, Li M (2011) Baicalein inhibits formation of α -synuclein oligomers within living cells and prevents A β peptide fibrillation and oligomerisation. *Chembiochem* **12**, 615-624.
- [63] Gertz M, Nguyen GT, Fischer F, Suenkel B, Schlicker C, Fränzel B, Tomaschewski J, Aladini F, Becker C, Wolters D, Steegborn C (2012) A molecular mechanism for direct sirtuin activation by resveratrol. *PLoS One* **7**, e49761.
- [64] Loh KP, Qi J, Tan BK, Liu XH, Wei BG, Zhu YZ (2010) Leonurine protects middle cerebral artery occluded rats through antioxidant effect and regulation of mitochondrial function. *Stroke* **41**, 2661-2668.
- [65] Shi XR, Hong ZY, Liu HR, Zhang YC, Zhu YZ (2011) Neuroprotective effects of SCM198 on 6-hydroxydopamine-induced behavioral deficit in rats and cytotoxicity in neuronal SH-SY5Y cells. *Neurochem Int* **58**, 851-860.
- [66] Hong ZY, Yu SS, Wang ZJ, Zhu YZ (2015) SCM-198 ameliorates cognitive deficits, promotes neuronal survival and enhances CREB/BDNF/TrkB signaling without affecting A β burden in A β PP/PS1 mice. *Int J Mol Sci* **16**, 18544-18563.
- [67] Jang SI, Pae HO, Choi BM, Oh GS, Jeong S, Lee HJ, Kim HY, Kang KJ, Yun YG, Kim YC, Chung HT (2003) Salidroside from *Rhodiola sachalinensis* protects neuronal PC12 cells against cytotoxicity induced by amyloid-beta. *Immunopharmacol Immunotoxicol* **25**, 295-304.
- [68] Zhang L, Yu H, Zhao X, Lin X, Tan C, Cao G, Wang Z (2010) Neuroprotective effects of salidroside against beta-amyloid-induced oxidative stress in SH-SY5Y human neuroblastoma cells. *Neurochem Int* **57**, 547-555.
- [69] Qu ZQ, Zhou Y, Zeng YS, Lin YK, Li Y, Zhong ZQ, Chan WY (2012) Protective effects of a *Rhodiola crenulata* extract and salidroside on hippocampal neurogenesis against streptozotocin-induced neural injury in the rat. *PLoS One* **7**, e29641.
- [70] Yang F, Lim GP, Begum AN, Ubeda OJ, Simmons MR, Ambegaokar SS, Chen PP, Kaye R, Glabe CG, Frautschy SA, Cole GM (2005) Curcumin inhibits formation of amyloid beta oligomers and fibrils, binds plaques, and reduces amyloid in vivo. *J Biol Chem* **280**, 5892-5901.
- [71] Ho YS, So KF, Chang RC (2011) Drug discovery from Chinese medicine against neurodegeneration in Alzheimer's and vascular dementia. *Chin Med* **6**, 15.
- [72] Wang R, Tang XC (2005) Neuroprotective effects of huperzine A. A natural cholinesterase inhibitor for the treatment of Alzheimer's disease. *Neurosignals* **14**, 71-82.
- [73] Zhu N, Lin J, Wang K, Wei M, Chen Q, Wang Y (2015) Huperzine A protects neural stem cells against A β -induced apoptosis in a neural stem cells and microglia co-culture system. *Int J Clin Exp Pathol* **8**, 6425-6433.
- [74] Ved HS, Koenig ML, Dave JR, Doctor BP (1997) Huperzine A, a potential therapeutic agent for dementia, reduces neuronal cell death caused by glutamate. *Neuroreport* **8**, 963-968.
- [75] Kim MK, Choi WY, Lee HY (2015) Enhancement of the neuroprotective activity of *Hericium erinaceus* mycelium co-cultivated with *Allium sativum* extract. *Arch Physiol Biochem* **121**, 19-25.
- [76] Muthukumaran V, Govindaraju M, Mohandass S, Visvanathan P, Kumar RS (2008) Evaluation of the neuroprotective role of Garlic (*Allium sativum*) extract and Methionine on mercury induced toxicity-an in vitro study. *Current Biotica* **2**, 161-172.
- [77] Gu X, Zhu YZ (2011) Therapeutic applications of organosulfur compounds as novel hydrogen sulfide donors and/or mediators. *Expert Rev Clin Pharmacol* **4**, 123-133.
- [78] Gong QH, Wang Q, Pan LL, Liu XH, Xin H, Zhu YZ (2011) S-propargyl-cysteine, a novel hydrogen sulfide-modulated agent, attenuates lipopolysaccharide-induced spatial learning and memory impairment: Involvement of TNF signaling and NF- κ B pathway in rats. *Brain Behav Immun* **25**, 110-119.
- [79] Birks J (2006) Cholinesterase inhibitors for Alzheimer's disease. *Cochrane Database Syst Rev*. CD005593.
- [80] Farlow M (2002) A clinical overview of cholinesterase inhibitors in Alzheimer's disease. *Int Psychogeriatr* **14**(Suppl 1), 93-126.
- [81] Zec RF, Burkett NR (2008) Non-pharmacological and pharmacological treatment of the cognitive and behavioral symptoms of Alzheimer disease. *NeuroRehabilitation* **23**, 425-438.
- [82] Ballard C, Corbett A (2010) Management of neuropsychiatric symptoms in people with dementia. *CNS Drugs* **24**, 729-739.
- [83] Wong GT, Manfra D, Poulet FM, Zhang Q, Josien H, Bara T, Engstrom L, Pinzon-Ortiz M, Fine JS, Lee HJ, Zhang L, Higgins GA, Parker EM (2004) Chronic treatment with the gamma-secretase inhibitor LY-411,575 inhibits beta-amyloid peptide production and alters lymphopoiesis and intestinal cell differentiation. *J Biol Chem* **279**, 12876-12882.
- [84] Gura T (2008) Hope in Alzheimer's fight emerges from unexpected places. *Nat Med* **14**, 894.
- [85] Martinez A, Perez DI (2008) GSK-3 inhibitors: A ray of hope for the treatment of Alzheimer's disease? *J Alzheimers Dis* **15**, 181-191.
- [86] Jankovic J (2005) Motor fluctuations and dyskinesias in Parkinson's disease: Clinical manifestations. *Mov Disord* **20**(Suppl 11), S11-S16.
- [87] Jankovic J, Aguilar LG (2008) Current approaches to the treatment of Parkinson's disease. *Neuropsychiatr Dis Treat* **4**, 743-757.

- [88] Li QY, Feng NH, Liu W (2011) Analysis of problems in the clinical application of Chinese medicine. *Western J Tradit Chin Med* **24**, 25-27.
- [89] Ferrer I (2012) Defining Alzheimer as a common age-related neurodegenerative process not inevitably leading to dementia. *Prog Neurobiol* **97**, 38-51.
- [90] Epstein LJ (1976) Depression in the elderly. *J Gerontol* **31**, 278-282.
- [91] Huang HJ, Chen HY, Lee CC, Chen CY (2014) Computational design of apolipoprotein E4 inhibitors for Alzheimer's disease therapy from traditional Chinese medicine. *Biomed Res Int* **2014**, 452625.
- [92] Suo T, Wang H, Li Z (2016) Application of proteomics in research on traditional Chinese medicine. *Expert Rev Proteomics* **13**, 873-881.
- [93] Wong VK, Law BY, Yao XJ, Chen X, Xu SW, Liu L, Leung EL (2016) Advanced research technology for discovery of new effective compounds from Chinese herbal medicine and their molecular targets. *Pharmacol Res* **111**, 546-555.